

Existence and uniqueness for planar anisotropic and crystalline curvature flow

A. Chambolle, M. Novaga

Abstract.

We prove short-time existence of φ -regular solutions to the planar anisotropic curvature flow, including the crystalline case, with an additional forcing term possibly unbounded and discontinuous in time, such as for instance a white noise. We also prove uniqueness of such solutions when the anisotropy is smooth and elliptic. The main tools are the use of an implicit variational scheme in order to define the evolution, and the approximation with flows corresponding to regular anisotropies.

CONTENTS

1. Introduction	1
2. RW_φ -condition and φ -regular flows	2
3. The time-discrete implicit scheme	6
4. Smooth anisotropies	8
5. General anisotropies	14

§1. Introduction

In this paper we consider the anisotropic curvature flow of planar curves, corresponding to the evolution law

$$(1) \quad V = \kappa_\varphi + \frac{\partial G}{\partial t}$$

in the Cahn-Hoffmann direction n_φ . We shall assume that the forcing term G has the form $G = G_1 + G_2$ with G_1, G_2 satisfying:

- i) $G_1 \in C^0([0, \infty))$ does not depend on x ;
- ii) $G_2 \in C^1([0, \infty); \text{Lip}(\mathbb{R}^2))$.

Observe that (1) is only formal, as $\partial G_1 / \partial t$ does not necessarily exist, however the motion can still be defined in an appropriate way (see Definition 2). Notice also that we include the case of G being a typical path of a Brownian motion, which is necessary to take into account a stochastic forcing term as in [18, 26].

In the smooth anisotropic case, the first existence and uniqueness results of a classical evolution in can be found in [5], where S. Angenent showed existence, uniqueness and comparison for a class of equations which include (1) in the case $G = G_2$ and φ regular. The existence and uniqueness of a weak solution for the forced flow, with a Lipschitz continuous forcing term, follows from standard viscosity theory [16, 17].

The crystalline curvature flow was mathematically formalized by J. Taylor in a series of papers (see for instance [28, 29]). In two-dimensions, the existence of the flow in the non forced case $G = 0$

Received Month Day, Year.

Revised Month Day, Year.

2000 *Mathematics Subject Classification*. Primary 53C44, 74N05, 74E10; Secondary: 35K55.

Key words and phrases. Anisotropy, Implicit variational scheme, Geometric evolutions, Crystal growth.

reduces to the analysis of a system of ODEs. It was first shown by F.J. Almgren and J. Taylor in [1], together with a proof of consistency of a variational scheme similar to the one introduced in Section 3. The uniqueness and comparison principle in the non forced case were established shortly after by Y. Giga and M.E. Gurtin in [25]. The forced crystalline flow was studied in [8], however with strong hypotheses on the forcing to ensure the preservation of the facets.

It is only in relatively recent work that the flow has been studied with quite general forcing terms: in [23, 24] a Lipschitz forcing is considered. However, [23] is restricted to the evolution of graphs (although with a general mobility), while [24] only considers rectangular anisotropies, and assumes that the initial datum is close to the Wulff shape. The paper [15] deals with quite general forcing terms (roughly, the same as in this paper), but requires the anisotropy to be smooth. It shows the consistency of the variational scheme and a comparison for regular evolutions. In [7], the authors show the existence of convex crystalline evolutions (extending their results of [6]) with time-dependent (bounded) forcing terms and apply it to show the existence of volume preserving flows.

We show here a general existence result for the two-dimensional crystalline curvature flow, with “natural” mobility, which holds in two cases: for a general forcing $G = G_1$ depending only on time, and for a regular forcing $G = G_2$ with $\partial G_2 / \partial t$ continuous in time and Lipschitz continuous in space.

Our proof relies on estimates for the variational scheme introduced in [2, 27], which show that, if the initial curve has a strong regularity (expressed in terms of an internal and external Wulff shape condition), then this regularity is preserved for some time which depends only on the initial radius. Extending these proofs to higher dimension would require quite strong regularity results for nonlinear elliptic PDEs, which do not seem available at a first glance.

The paper is organized as follows: in the Section 2 we define the “anisotropy” and introduce our notion of a “regular” curvature flow for smooth and nonsmooth anisotropies. In Section 3 we study the time-discrete implicit scheme of [2], and extend some regularity results of [6] to the flow with forcing. We then show in Section 4 the main existence results, for smooth anisotropies. The fundamental point is that the time of existence only depends on the regularity of the initial curve. In the smooth case, we also show uniqueness of regular evolutions. Eventually, in Section 5 we extend the existence result to the crystalline case, however this simply follows from an elementary approximation lemma (Lemma 1), and the fact that the time of existence is uniformly controlled in this approximation.

§2. RW_φ -condition and φ -regular flows

We call *anisotropy* a function φ which is convex, one-homogeneous and coercive on \mathbb{R}^2 . We will also assume that φ is even, i.e. φ is a norm, although we expect that the results of this paper still hold in the general case (but some proofs become more tedious to write).

We will always assume that there exists $c_0 > 0$ such that

$$(2) \quad c_0 |x| \leq \varphi(x) \leq c_0^{-1} |x| \quad \forall x \in \mathbb{R}^2.$$

We denote by φ° the polar of φ , defined as

$$\varphi^\circ(\nu) := \sup_{x: \varphi(x) \leq 1} \nu \cdot x \quad \nu \in \mathbb{R}^2,$$

it is obviously also a convex, one-homogeneous and even function on \mathbb{R}^2 . Notice that from (2) it easily follows

$$c_0 |\nu| \leq \varphi^\circ(\nu) \leq c_0^{-1} |\nu| \quad \forall \nu \in \mathbb{R}^2.$$

We denote by $W_\varphi := \{\varphi \leq 1\}$ the unit ball of φ , which is usually called the *Wulff shape*.

We say that φ is *smooth* if $\varphi \in C^2(\mathbb{R}^2 \setminus \{0\})$ and φ is *elliptic* if φ^2 is strictly convex, that is $\nabla^2(\varphi^2) \geq c \text{Id}$ in the distributional sense, for some $c > 0$. It is easy to check that φ is smooth and elliptic iff φ° is smooth and elliptic.




Fig. 1. The pathological set described in Remark 1.

Given a set $E \subset \mathbb{R}^2$ we let d_φ^E be the signed φ -distance function to ∂E defined as

$$d_\varphi^E(x) := \min_{y \in E} \varphi(x - y) - \min_{y \notin E} \varphi(y - x),$$

We let $\nu_\varphi^E := \nabla d_\varphi^E$ be the exterior φ -normal to ∂E , $n_\varphi \in \partial\varphi^\circ(\nu_\varphi)$ be the so-called *Cahn-Hoffmann* vector field (where ∂ denotes the subdifferential), and $\kappa_\varphi := \operatorname{div} n_\varphi$ be the φ -curvature of ∂E , whenever they are defined. We also set $E^c := \mathbb{R}^2 \setminus E$.

Following [11] we give the following definition:

Definition 1 (*RW_φ -condition*). We say that a set E satisfies the inner RW_φ -condition for some $R > 0$ if

$$(3) \quad \overline{E} = \bigcup_{x: d_\varphi^E(x) \leq -R} (x + RW_\varphi)$$

and for any $r < R$ and $x \in \mathbb{R}^2$, $(x + rW_\varphi) \cap E^c$ is connected.

We say that E satisfies the outer RW_φ -condition if its complementary E^c satisfies the inner RW_φ -condition.

We say that E satisfies the RW_φ -condition if it satisfies both the inner and outer RW_φ -conditions.

Remark 1. Notice that, if E satisfies the RW_φ -condition for some $R > 0$, then ∂E is locally a Lipschitz graph. Moreover, when φ is smooth, the RW_φ -condition implies that ∂E is of class $C^{1,1}$ and $|\kappa_\varphi| \leq 1/R$ a.e. on ∂E . In this case the connectedness condition in Definition 1 is automatically satisfied whenever (3) holds. However, in the nonsmooth case one can have some pathological examples if one removes the connectedness condition, as the one depicted in Fig. 1 when the Wulff shape is a square.

Remark 2. It is not difficult to show that E satisfies the inner RW_φ -condition iff (3) holds and the following property holds: for all x such that $d_\varphi^E(x) = -R' > -R$, the set $\partial E \cap (x + R'\partial W_\varphi)$ is connected.

By (3) it follows that $\partial E \cap (x + R'\partial W_\varphi)$ is either a segment (possibly a point) or the union of two segments. In particular, if φ is elliptic, this is equivalent to say that there exists a unique point in ∂E minimizing the φ -distance from x .

Definition 2 (*φ -regular flows*). We say that a map $[0, T] \ni t \rightarrow \mathcal{P}(\mathbb{R}^2)$ defines a φ -regular flow for (1) if

- (1) $E(t)$ satisfies the RW_φ -condition for all $t \in [0, T]$ and for some $R > 0$;
- (2) there exist an open set $U \subset \mathbb{R}^2$ and a vector field $z \in L^\infty([0, T] \times U; \mathbb{R}^2)$ such that

- (a) $\partial E(t) \subset U$ for all $t \in [0, T]$,
 - (b) $d_\varphi^E(t, x) := d_\varphi^{E(t)}(x) \in C^0([0, T]; \text{Lip}(U))$,
 - (c) $z \in \partial\varphi^\circ(\nabla d_\varphi^E)$ a.e. in $[0, T] \times U$,
 - (d) $\text{div} z \in L^\infty([0, T] \times U)$;
- (3) there exists $\lambda > 0$ such that, for any t, s with $0 \leq t < s \leq T$ and a.e. $x \in U$, there holds

$$(4) \quad \left| d_\varphi^E(s, x) - d_\varphi^E(t, x) - \int_s^t \text{div} z(\tau, x) d\tau - G(s, x) + G(t, x) \right| \leq \lambda(s - t) \max_{t \leq \tau \leq s} |d_\varphi^E(x, \tau)|.$$

Observe that (4) implies that $(d - G)$ is Lipschitz continuous, so that (4) can be rewritten as

$$(5) \quad \left| \frac{\partial(d_\varphi^E - G)}{\partial t}(t, x) - \text{div} \nabla\varphi^\circ(\nabla d_\varphi^E)(t, x) \right| \leq \lambda |d_\varphi^E(t, x)|.$$

for a.e. $(t, x) \in [0, T] \times U$. In case G is C^1 in time, equation (5) expresses the fact that $\partial E(t)$ evolves with speed given by (1).

2.1. An approximation result.

We now show that, given any set E satisfying the RW_φ -condition for a general anisotropy φ , there exist smooth and elliptic anisotropies $\varphi_\varepsilon \rightarrow \varphi$ and sets $E_\varepsilon \rightarrow E$, as $\varepsilon \rightarrow 0$, such that E_ε satisfies the RW_{φ_ε} -condition.

Lemma 1. *Let φ be a general anisotropy and let φ_ε be smooth and elliptic anisotropies converging to φ , with $\varphi_\varepsilon \geq \varphi$. Let $E \subseteq \mathbb{R}^2$ satisfy the RW_φ -condition for some $R > 0$. Then there exist sets E_ε , with $\partial E_\varepsilon \rightarrow \partial E$ as $\varepsilon \rightarrow 0$ in the Hausdorff sense, such that each E_ε satisfies the RW_{φ_ε} -condition.*

Proof. Let

$$\begin{aligned} \tilde{E}_\varepsilon &:= \bigcup \{ (x + RW_{\varphi_\varepsilon}) : (x + RW_{\varphi_\varepsilon}) \subset \overline{E} \} \\ E_\varepsilon &:= \mathbb{R}^2 \setminus \bigcup \left\{ (x + RW_{\varphi_\varepsilon}) : (x + RW_{\varphi_\varepsilon}) \subset \overline{\tilde{E}_\varepsilon^c} \right\}. \end{aligned}$$

Notice that, by definition, \tilde{E}_ε satisfies the inner RW_{φ_ε} -condition and E_ε satisfies the outer RW_{φ_ε} -condition, so that we have to prove that E_ε also satisfies the inner RW_{φ_ε} -condition.

Step 1. Let us show that $\partial E_\varepsilon \rightarrow \partial E$ as $\varepsilon \rightarrow 0$, in the Hausdorff sense. In fact, this is obvious from the construction: since $W_{\varphi_\varepsilon} \subset W_\varphi$ and for any $x \in E$, there exists $z \in E$ with $x \in z + RW_\varphi \subset E$, we see that the distance from x to \tilde{E}_ε (and then E_ε) is bounded by the Hausdorff distance between RW_φ and RW_{φ_ε} . An estimate for the complement can be derived in the same way, so that $d_{\mathcal{H}}(\partial E_\varepsilon, \partial E) \leq R d_{\mathcal{H}}(W_\varphi, W_{\varphi_\varepsilon})$.

Step 2. We now prove that \tilde{E}_ε satisfies the outer RW_φ -condition. We first show that, for all $x \in \partial \tilde{E}_\varepsilon$, there exists y such that

$$(6) \quad \tilde{E}_\varepsilon \subset (y + RW_\varphi)^c \quad \text{and} \quad x \in \partial(y + RW_\varphi).$$

Indeed, if $x \in \partial \tilde{E}_\varepsilon \cap \partial E$, (6) readily follows from the fact that E satisfies the outer RW_φ -condition.

If $x \in \partial \tilde{E}_\varepsilon \setminus \partial E$, then by definition of \tilde{E}_ε there exists $x_1 \in \mathbb{R}^2$ such that

$$(x_1 + RW_{\varphi_\varepsilon}) \subseteq \tilde{E}_\varepsilon \quad \text{and} \quad x \in \partial(x_1 + RW_{\varphi_\varepsilon}).$$

Let ℓ_x be the maximal arc of $\partial(x_1 + RW_{\varphi_\varepsilon})$ containing x and contained in the interior of E , and let $y_1, y_2 \in \partial E$ be the endpoints of ℓ_x . Notice that $\varphi_\varepsilon(y_1 - y_2) < 2R$. Let $y_3 := (y_1 + y_2)/2$, $R' := \varphi(y_1 - y_2)/2 \leq \varphi_\varepsilon(y_1 - y_2)/2 < R$. As E satisfies the inner RW_φ -condition, the set $(y_3 + R'W_\varphi)$ has connected intersection with E^c , so that the set $(\partial E \setminus \partial \tilde{E}_\varepsilon) \cap (y_3 + R'W_\varphi)$ contains a connected arc $\tilde{\ell}_x$ joining y_1 and y_2 (see Figure 2).




Fig. 2. The configuration in Lemma 1.

Let S_x be the subset of E such that $\partial S_x = \ell_x \cup \tilde{\ell}_x$ and set $E' := \tilde{E}_\varepsilon \cap (y_3 + R'W_\varphi)$. Notice that $S_x \subset E'$. As E' is a convex set, there exists y such that $x \in \partial(y + RW_\varphi)$ and $E' \subset (y + RW_\varphi)^c$. Moreover, since E satisfies the outer RW_φ -condition, the set $E \cap \text{int}(y + RW_\varphi) \supseteq S_x \cap \text{int}(y + RW_\varphi)$ is connected. This implies that $\tilde{E}_\varepsilon \subset (y + RW_\varphi)^c$ and proves (6).

In order to prove that \tilde{E}_ε satisfies the outer RW_φ -condition, by Remark 2 it remains to show that, given \bar{x} with $d_{\varphi_\varepsilon}^{\tilde{E}_\varepsilon} = R' < R$, the set $\partial \tilde{E}_\varepsilon \cap (\bar{x} + R'\partial W_\varphi)$ is connected.

If $\partial \tilde{E}_\varepsilon \cap (\bar{x} + R'\partial W_\varphi) \subset \partial E$ this follows directly from the fact that E satisfies the outer RW_φ -condition. Otherwise, there exists $x \in (\partial \tilde{E}_\varepsilon \setminus \partial E) \cap (\bar{x} + R'\partial W_\varphi)$. In this case we claim that $\partial \tilde{E}_\varepsilon \cap (\bar{x} + R'\partial W_\varphi) = \{x\}$. Indeed, since ℓ_x is a strictly convex arc, we have $\ell_x \cap (\bar{x} + R'\partial W_\varphi) = \{x\}$. Hence, if $\partial \tilde{E}_\varepsilon \cap (\bar{x} + R'\partial W_\varphi)$ contains another point $y \neq x$, then $y \notin \overline{S_x}$. As $S_x \cap (\bar{x} + R'W_\varphi) \neq \emptyset$, it follows that $E \cap (\bar{x} + (R' + \delta)W_\varphi)$ contains at least two connected components for $\delta > 0$ sufficiently small, contradicting the fact that E satisfies the outer RW_φ -condition. Hence the set $(\partial E \setminus \partial \tilde{E}_\varepsilon) \cap (y_3 + R'W_{\varphi_\varepsilon})$ is a connected arc $\tilde{\ell}_x$ joining y_1 and y_2 .

Step 3. We prove that E_ε satisfies the inner RW_{φ_ε} -condition by reasoning as in *Step 2*, with E replaced by $(\tilde{E}_\varepsilon)^c$ (and φ replaced by φ_ε). The only difference is due to the fact that $(\tilde{E}_\varepsilon)^c$ now satisfies inner RW_φ -condition and the outer RW_{φ_ε} -condition. Therefore, letting $R' := \varphi_\varepsilon(y_1 - y_2)/2 < R$, the set $(y_3 + R'W_\varphi) \cap \tilde{E}_\varepsilon$ is connected, so that $(\partial \tilde{E}_\varepsilon \setminus \partial E_\varepsilon) \cap (y_3 + R'W_{\varphi_\varepsilon})$ contains a connected arc joining y_1 and y_2 . In the rest of the proof one can proceed as in *Step 2*.

Q.E.D.

Lemma 1 has the following direct consequence.

Corollary 1. *Let $E \subseteq \mathbb{R}^2$ satisfy the RW_φ -condition for some $R > 0$. Then E is φ -regular in the sense of [11], that is, there exists a vector field $n_\varphi \in L^\infty(\{|d_\varphi^E| < R\}, \mathbb{R}^2)$ such that $n_\varphi \in \partial \varphi^\circ(\nabla d_\varphi^E)$ a.e. in $\{|d_\varphi^E| < R\}$, and $\text{div} n_\varphi \in L_{\text{loc}}^\infty(\{|d_\varphi^E| < R\})$.*

Proof. Take a sequence φ_ε of smooth and elliptic anisotropies converging to φ , with $\varphi_\varepsilon \geq \varphi$. By Lemma 1 we can approximate E in the Hausdorff distance with sets E_ε satisfying the RW_{φ_ε} -condition. In particular, letting $n_{\varphi_\varepsilon} = \nabla \varphi_\varepsilon^\circ(\nabla d_{\varphi_\varepsilon}^{E_\varepsilon}) \in L^\infty(\mathbb{R}^2)$ and recalling Remark 1, we have that $\text{div} n_{\varphi_\varepsilon} \in L_{\text{loc}}^\infty(\{|d_{\varphi_\varepsilon}^{E_\varepsilon}| < R\})$. Therefore, any weak* limit n_φ of n_{φ_ε} , as $\varepsilon \rightarrow 0$, satisfies the thesis.

Q.E.D.

Remark 3. Notice that, given an arbitrary anisotropy φ , it is relatively easy to approximate it with smooth and elliptic anisotropies φ_ε . For instance, one may let $F_\varepsilon := \{\eta_\varepsilon * \varphi^\circ \leq 1\} \oplus B(0, \varepsilon)$,

with $\eta_r(x) := r^{-d}\eta(\frac{x}{r})$, and $\varphi_\varepsilon(x) := \sup_{\nu \in F_\varepsilon} \nu \cdot x$. It is easy to check that the anisotropies φ_ε are smooth and elliptic, and converge locally uniformly to φ as $\varepsilon \rightarrow 0$.

§3. The time-discrete implicit scheme

The results of this section hold in any dimension $d \geq 2$ and are stated in this general form. Up to minor improvements, they are essentially stated in [15, 6]. Following [15] we recall the definition and some properties of the implicit scheme introduced in [2, 27]. Given a set $E \subset \mathbb{R}^d$ with compact boundary (we assume without loss of generality that it is bounded), we define for $s > t \geq 0$ a transformation $T_{t,s}$ by letting $T_{t,s}(E) = \{x \in B_R : w(x) < 0\}$, where $B_R = B(0, R)$, R is large and w is the minimizer of

$$(7) \quad \min_{w \in L^2(B_R)} \int_{B_R} \varphi^\circ(Dw) + \frac{1}{2(s-t)} \int_{B_R} \left(w(x) - d_\varphi^E(x) - G(s, x) + G(t, x) \right)^2 dx,$$

whose existence and uniqueness is shown by standard methods. One checks easily [13, 15, 3] that for R large, the level set $T_{t,s}(E)$ of w does not depend on R , and it is a solution to the variational problem

$$(8) \quad \min P_\varphi(F) + \frac{1}{s-t} \int_F \left(d_\varphi^E(x) + G(s, x) - G(t, x) \right) dx,$$

where the minimum is taken among the subsets F of \mathbb{R}^d with finite perimeter, and we set

$$P_\varphi(F) := \int_{\partial^* F} \varphi^\circ(\nu_F(x)) d\mathcal{H}^1(x).$$

It follows that the set $T_{t,s}(E)$ has boundary of class $C^{1,\alpha}$, outside a compact singular set of zero \mathcal{H}^1 -dimension [2] (when $d = 2$, the set $T_{t,s}(E)$ has boundary of class $C^{1,1}$). The variational problem above is the generalization of the approach proposed in [2, 27], for building mean curvature flows without driving terms, through an implicit time discretization.

For $s = t + h$, the Euler-Lagrange equation for $T_{t,t+h}(E)$ at a point $x \in \partial T_{t,t+h}(E)$ formally reads as

$$d_\varphi^E(x) = -h \left(\kappa_\varphi(x) + \frac{G(t+h, x) - G(t, x)}{h} \right),$$

with κ_φ being the φ -curvature at x of $\partial T_{t,t+h}(E)$, so that it corresponds to an implicit time-discretization of (1). Observe also that this approximation is trivially monotone: indeed if $E \subseteq E'$ then $d_\varphi^E \geq d_\varphi^{E'}$, which yields $w \geq w'$, w and w' being the solutions of (7) for the distance functions d_φ^E and $d_\varphi^{E'}$ respectively. We deduce that $\{w < 0\} \subseteq \{w' < 0\}$, that is,

$$(9) \quad E \subseteq E' \implies T_{t,t+h}(E) \subseteq T_{t,t+h}(E').$$

Consider now the Euler-Lagrange equation for (7), which is

$$(10) \quad -(s-t)\operatorname{div} z + w(x) = d_\varphi^E(x) + G(s, x) - G(t, x)$$

for $x \in B_R$, with $\varphi(z(x)) \leq 1$ and $z(x) \cdot \nabla w(x) = \varphi^\circ(\nabla w(x))$ a.e. in B_R (by elliptic regularity one knows that w is Lipschitz).

We show that if E is regular enough, then we have an estimate on the quantity $\operatorname{div} z + (G(s, x) - G(t, x))/(s-t)$ near the boundary of E . The technique is adapted from [6].

Lemma 2. *Assume that E is a bounded set which satisfies the $\delta\mathcal{W}_\varphi$ -condition for some $\delta > 0$. Let $a < b$ be such that $X_{a,b} := \{\max\{w, d_\varphi^E\} \geq a\} \cap \{\min\{w, d_\varphi^E\} \leq b\} \subseteq \{|d_\varphi^E| < \delta\}$. Then $\operatorname{div} z \in L^\infty(X_{a,b})$ and*

$$(11) \quad \left\| \operatorname{div} z + \frac{G(s, \cdot) - G(t, \cdot)}{s-t} \right\|_{L^\infty(X_{a,b})} \leq \left\| \operatorname{div} n_\varphi^E + \frac{G(s, \cdot) - G(t, \cdot)}{s-t} \right\|_{L^\infty(X_{a,b})}.$$

Proof. Let $f : \mathbb{R} \rightarrow [0, +\infty)$ be a smooth increasing function with $f(t) = 0$ if $t \leq 0$. Since (w, z) solves (10), we find

$$\begin{aligned} \int_{X_{a,b}} (w - d_\varphi^E) f(w - d_\varphi^E) dx &= \int_{X_{a,b}} ((s-t) \operatorname{div} z + G(s, x) - G(t, x)) f(w - d_\varphi^E) dx \\ &= (s-t) \int_{X_{a,b}} (\operatorname{div} z - \operatorname{div} n_\varphi^E) f(w - d_\varphi^E) dx \\ &\quad + \int_{X_{a,b}} ((s-t) \operatorname{div} n_\varphi^E + G(s, x) - G(t, x)) f(w - d_\varphi^E) dx =: \text{I} + \text{II}. \end{aligned}$$

We have, observing that $X_{a,b}$ has Lipschitz boundary (for a.e. choice of a, b),

$$\begin{aligned} \text{I} &= - (s-t) \int_{X_{a,b}} (z - n_\varphi^E) \cdot \nabla (w - d_\varphi^E) f'(w - d_\varphi^E) dx \\ &\quad + (s-t) \int_{\partial X_{a,b}} f(w - d_\varphi^E) (z - n_\varphi^E) \cdot \nu^{X_{a,b}} d\mathcal{H}^1 =: \text{I}_1 + \text{I}_2. \end{aligned}$$

First $\text{I}_1 \leq 0$ since $z \cdot \nabla w = \varphi^\circ(\nabla w)$ and $z \cdot \nabla d_\varphi^E = \varphi^\circ(\nabla d_\varphi^E)$. We claim that also $\text{I}_2 \leq 0$. Indeed, on one hand, when $f(w - d_\varphi^E) > 0$, $w > d_\varphi^E$ hence, $\nu^{X_{a,b}} = \nu^{\{d_\varphi^E \leq b\}} = \nabla d_\varphi^E / |\nabla d_\varphi^E|$, \mathcal{H}^1 -almost everywhere on $\{\min\{w, d_\varphi^E\} = b\}$ while $\nu^{X_{a,b}} = \nu^{\{w \geq a\}} = -\nabla w / |\nabla w|$, \mathcal{H}^1 -almost everywhere on $\{\max\{w, d_\varphi^E\} = a\}$. It follows that $f(w - d_\varphi^E)(z - n_\varphi^E) \cdot \nu^{X_{a,b}} \leq 0$ on both $\{\min\{w, d_\varphi^E\} = b\}$ and $\{\max\{w, d_\varphi^E\} = a\}$ so that $\text{I}_2 \leq 0$. We conclude that $\text{I} \leq 0$, hence

$$(12) \quad \int_{X_{a,b}} (w - d_\varphi^E) f(w - d_\varphi^E) dx \leq \int_{X_{a,b}} ((s-t) \operatorname{div} n_\varphi^E + G(s, x) - G(t, x)) f(w - d_\varphi^E) dx.$$

Let $q > 2$, let $r^+ := r \vee 0$, and let $\{f_n\}$ be a sequence of smooth increasing nonnegative functions such that $f_n(r) \rightarrow r^{+(q-1)}$ uniformly as $n \rightarrow \infty$. From (12) we obtain

$$\begin{aligned} \int_{X_{a,b}} ((w - d_\varphi^E)^+)^q dx &\leq \int_{X_{a,b}} ((s-t) \operatorname{div} n_\varphi^E + G(s, x) - G(t, x)) ((w - d_\varphi^E)^+)^{q-1} dx \\ &\leq \int_{X_{a,b}} ((s-t) \operatorname{div} n_\varphi^E + G(s, x) - G(t, x))^+ ((w - d_\varphi^E)^+)^{q-1} dx. \end{aligned}$$

Applying Young's inequality we obtain

$$\|(w - d_\varphi^E)^+\|_{L^q(X_{a,b})} \leq \left\| ((s-t) \operatorname{div} n_\varphi^E + G(s, \cdot) - G(t, \cdot))^+ \right\|_{L^q(X_{a,b} \cap \{w > d_\varphi^E\})}.$$

A similar proof, reverting the signs, shows that

$$\|(w - d_\varphi^E)^-\|_{L^q(X_{a,b})} \leq \left\| ((s-t) \operatorname{div} n_\varphi^E + G(s, \cdot) - G(t, \cdot))^- \right\|_{L^q(X_{a,b} \cap \{w < d_\varphi^E\})}.$$

It follows that

$$\|(s-t) \operatorname{div} z + G(s, \cdot) - G(t, \cdot)\|_{L^q(X_{a,b})} \leq \|(s-t) \operatorname{div} n_\varphi^E + G(s, \cdot) - G(t, \cdot)\|_{L^q(X_{a,b})},$$

and letting $q \rightarrow \infty$ we obtain (11). Observe that the estimate we may obtain is a bit more precise, in fact we have shown:

$$\begin{aligned} (13) \quad \operatorname{ess\,inf}_{X_{a,b} \cap \{w < d_\varphi^E\}} \operatorname{div} n_\varphi^E + \frac{G(s, \cdot) - G(t, \cdot)}{s-t} \\ \leq \operatorname{div} z(x) + \frac{G(s, x) - G(t, x)}{s-t} \leq \operatorname{ess\,sup}_{X_{a,b} \cap \{w > d_\varphi^E\}} \operatorname{div} n_\varphi^E + \frac{G(s, \cdot) - G(t, \cdot)}{s-t} \end{aligned}$$

for a.e. $x \in X_{a,b}$.

Q.E.D.

We also recall Lemma 3.2 from [15]:

Lemma 3. *Let $x_0 \in B_R$ and $\rho > 0$, and let $t \geq 0$. Let \tilde{w} solve*

$$(14) \quad \min_{\tilde{w} \in L^2(B_R)} \int_{B_R} \varphi^\circ(D\tilde{w}) + \frac{1}{2h} \int_{B_R} (\tilde{w}(x) - (\varphi(x - x_0) - \rho) - G(x, t + h) + G(x, t))^2 dx.$$

Then

$$(15) \quad \tilde{w}(x) \leq \begin{cases} \varphi(x - x_0) + h \frac{1}{\varphi(x - x_0)} + \Delta_h(t) - \rho & \text{if } \varphi(x - x_0) \geq \sqrt{2h} \\ 2\sqrt{2h} + \Delta_h(t) - \rho & \text{otherwise,} \end{cases}$$

where $\Delta_h(t) := \|G(\cdot, t + h) - G(\cdot, t)\|_{L^\infty(B_R)}$.

We deduce an estimate on $w - d_\varphi^E$, if E has an inner ρW_φ -condition: indeed, in this case, if $h = s - t$,

$$d_\varphi^E(x) \leq \inf \{ \varphi(x - x_0) - \rho : d_\varphi^E(x_0) = -\rho \}$$

with, in fact, equality in $\{-\rho \leq d_\varphi^E \leq \rho'\}$, where $\rho' \geq 0$ is the radius of an outer $\rho' W_\varphi$ -condition. It follows from (15) that

$$(16) \quad w(x) \leq \inf \left\{ \varphi(x - x_0) - \rho + h \frac{1}{\varphi(x - x_0)} + \Delta_h(t) : d_\varphi^E(x_0) = -\rho \right\}$$

for x with $d_\varphi^E(x) \geq -\rho + \sqrt{2h}$, and more precisely if $\rho' \geq d_\varphi^E(x) \geq -\rho/2$,

$$(17) \quad w(x) \leq d_\varphi^E(x) + \frac{2h}{\rho} + \Delta_h(t),$$

as soon as $h \leq \rho^2/16$.

§4. Smooth anisotropies

4.1. Existence of φ -regular flows.

We will prove, in dimension $d = 2$, an existence result for the forced curvature flow, first in case the anisotropy is smooth and elliptic. For technical reason, we need the forcing term G to be either time-dependent only (case $G_2 = 0$), or smooth (globally Lipschitz in space and time, case $G_1 = 0$).

Theorem 1. *Assume $G_1 = 0$ or $G_2 = 0$, and let (φ, φ°) be a smooth and elliptic anisotropy and $E_0 \subset \mathbb{R}^2$ an initial set with compact boundary, satisfying both an RW_φ -internal and external condition. Then, there exist $T > 0$, and a φ -regular flow $E(t)$ defined on $[0, T]$ and starting from $E(0) = E_0$.*

More precisely, there exist $R' > 0$ and a neighborhood U of $\bigcup_{0 \leq t \leq T} \partial E(t)$ in \mathbb{R}^2 such that the sets $E(t)$ satisfy the $R' W_\varphi$ -condition for all $t \in [0, T]$, the φ -signed distance function $d_\varphi^E(t, x)$ from $\partial E(t)$ belongs to $C^0([0, T]; \text{Lip}(U)) \cap L^\infty([0, T]; C^{1,1}(U))$, $(d_\varphi^E - G) \in \text{Lip}([0, T] \times U)$ and

$$(18) \quad \left| \frac{\partial(d_\varphi^E - G)}{\partial t}(t, x) - \text{div } \nabla \varphi^\circ(\nabla d_\varphi^E)(t, x) \right| \leq \lambda |d_\varphi^E(t, x)|.$$

for a.e. $(t, x) \in [0, T] \times U$, where λ is a positive constant. Finally, the time T , the radius R' , the set U , and the constant λ depend only on R and G .

Theorem 1 will be proved by time-discretization. Before, we need the following technical lemma.

Lemma 4. *Let φ, φ° be smooth and elliptic, and a set E satisfy a RW_φ -condition for some $R > 0$. We also assume that E is simply connected (∂E is a $C^{1,1}$ Jordan curve). Let $\delta \in (0, R)$ and consider a set F (also simply connected), such that $E_\delta \subset F \subset E^\delta$. Assume that $\|\kappa_\varphi^F\|_{L^\infty(\partial F)} \leq K$ for a constant $K < 1/(2\delta)$. Then F has a $R'W_\varphi$ -condition, with $R' = \min\{R - \delta, (1 - 2\delta K)/K\}$.*

Proof. We assume that ∂F is at least C^2 . If the result holds in this case, then given a more general $C^{1,1}$ set we can smooth it slightly, use the result for the approximations, and then pass to the limit.

Step 1. We have $E^\delta \setminus E_\delta = \bigcup_{x \in \partial E} x + \delta W_\varphi$, and for any $x \in \partial E$, the set $x + \delta W_\varphi$ is tangent to ∂E_δ (respectively, ∂E^δ) at exactly one point $x - \delta n_\varphi(x)$ (resp., $x + \delta n_\varphi(x)$). We can define Γ_x^+ and Γ_x^- as the two arcs on $\partial(x + \delta W_\varphi)$ delimited by the points $x \pm \delta n_\varphi(x)$, the exponent $+$ and $-$ indicating that Γ_x^\pm meets ∂E right “after” or “before” x , relative to an arbitrarily chosen orientation of the curve.

A first observation is that $\sharp(\partial F \cap \Gamma_x^\pm) = 1$ for all x . Indeed, we check that this value is a continuous function of x . If not, there will exist for instance a point where $\sharp(\partial F \cap \Gamma_x^+)$ has a “jump”, that is, where ∂F is tangent to Γ_x^+ and contains a small piece of arc which is inside $x + \delta W_\varphi$ and tangent to its boundary: in this case, we deduce that $\kappa_\varphi^F(x)$ is larger than $1/\delta$ or less than $-1/\delta$, a contradiction.

Since this value is continuous, it can only be odd (since $E_\delta \subset F \subset E^\delta$), moreover if it were larger than 1, there would be a connected component of F (as well as one of its complement) in $E^\delta \setminus E_\delta$, a contradiction.

Step 2. Let $\rho < \min\{R - \delta, (1 - 2\delta K)/K\}$. Assume that there exists $y \in F$ such that $W := y + \rho W_\varphi \subset \overline{F}$ and $y + \rho W_\varphi$ meets ∂F in at least two points z^-, z^+ (with z^+ “after” z^- with respect to the orientation along ∂E). These points must be isolated (otherwise there would be a point on ∂F with curvature equal to $1/\rho > K$). Observe also that $W \cap (E^\delta \setminus E_\delta)$ is connected (since E^δ has an inner $(R - \delta)W_\varphi$ -condition). To z^+ , we can associate a unique x^+ such that $z^+ \in \Gamma_{x^+}^+$, and to z^- a unique x^- such that $z^- \in \Gamma_{x^-}^-$. Then, the piece of curve Γ of ∂F between z^- and z^+ lies in the region of $E^\delta \setminus E_\delta$ bounded by $\Gamma_{x^-}^-$ and $\Gamma_{x^+}^+$, which contains points at “distance” at most 2δ from W : more precisely, $\Gamma \subset y + (\rho + 2\delta)W_\varphi$. Hence, there exists $s \in (\rho, \rho + 2\delta]$ such that Γ is contained in $y + sW_\varphi$ and tangent to its boundary, and thus a point of curvature larger than $1/s \geq 1/(\rho + 2\delta) > K$ on ∂W_φ , which is a contradiction. Therefore, the Wulff shapes $y + \rho W_\varphi$ which lie inside F can touch its boundary at most in one point, and an inner condition of radius $\min\{R - \delta, (1 - 2\delta K)/K\}$ easily follows.

The proof of the outer condition is identical.

Q.E.D.

Remark 4. We can refine the lemma to consider a situation where E has an inner $R_i W_\varphi$ -condition and a outer $R_o W_\varphi$ -condition, for two given radii $R_i, R_o > \delta$. We assume that $-K_o \leq \kappa_\varphi^F \leq K_i$ for two nonnegative constants K_i, K_o , (still less than $1/(2\delta)$). It is then deduced that F has a inner $R'_i W_\varphi$ -condition and a outer $R'_o W_\varphi$ -condition, with $R'_i = \min\{R_i - \delta, (1 - 2\delta K_i)/K_i\}$, $R'_o = \min\{R_o - \delta, (1 - 2\delta K_o)/K_o\}$.

Proof of Theorem 1. From (11) and (13), we will obtain some regularity of the boundary of $T_{t+h,t}(E)$, which will allow to iterate the variational scheme. To simplify (and without loss of generality) we assume that the initial curve is a Jordan curve (E_0 is simply connected). If not, one may evolve separately each connected component of the boundary.

Step 1.a.: The case $G_1 = 0$. In the case $G_1 = 0$, there exists C such that $\Delta_h(t) \leq Ch$. It follows from (17) that if E satisfies the ρW_φ -condition, then, by solving (7) with $s = t + h$,

$$(19) \quad |w(x) - d_\varphi^E(x)| \leq h \left(C + \frac{2}{\rho} \right)$$

in $\{|d_\varphi^E| < \rho/2\}$. By standard comparison (using for instance Lemma 3 again) one also can check that $w < 0$ if $d_\varphi^E \leq \rho/2$, and $w > 0$ if $d_\varphi^E \geq \rho/2$, so that the boundary of $T_{t,t+h}(E)$ is at (φ) -distance of order h from ∂E , if $h \leq \rho^2/36$ (Lemma 3). We also observe that the Hausdorff distance between the sets E and $T_{t,t+h}(E)$ is of the same order, or equivalently, $\|d_\varphi^E - d_\varphi^{T_{t,t+h}(E)}\|_{L^\infty(\mathbb{R}^d)} \leq (C + 2/\rho)h$.

A further observation is that if E is simply connected, also $T_{t,t+h}(E)$ is. Indeed, if not, there would be a connected component of either $T_{t,t+h}(E)$ or its complement in the set $\{|d_\varphi^E| \leq h(C + 2/\rho)\}$. Assume F is a connected of $T_{t,t+h}(E)$ which lies in $\{|d_\varphi^E| \leq h(C + 2/\rho)\}$, so that $|F| \leq 2hP_\varphi(E)(C + 2/\rho)$. One has that (using the isoperimetric inequality)

$$\begin{aligned} P_\varphi(F) + \frac{1}{h} \int_F d_\varphi^E(x) + G(t+h, x) - G(t, x) dx \\ \geq 2\sqrt{|W_\varphi||F|} - 2|F| \left(C + \frac{1}{\rho}\right) \\ \geq 2\sqrt{|F|} \left(\sqrt{|W_\varphi|} - \sqrt{|F|} \left(C + \frac{1}{\rho}\right)\right) \end{aligned}$$

which is positive if h is small enough (depending on $C, \rho, P_\varphi(E)$), showing that $T_{t,t+h}(E) \setminus F$ has an energy strictly lower than $T_{t,t+h}(E)$ in (8), a contradiction.

Sending both a and b to 0, one deduces from (11) that $T_{t,t+h}(E)$ has $C^{1,1}$ boundary, and moreover

$$\begin{aligned} \left\| \operatorname{div} n_\varphi^{T_{t,t+h}(E)} + \frac{1}{h}(G(t+h, \cdot) - G(t, \cdot)) \right\|_{L^\infty(\partial T_{t,t+h}(E))} \\ \leq \left\| \operatorname{div} n_\varphi^E + \frac{1}{h}(G(t+h, \cdot) - G(t, \cdot)) \right\|_{L^\infty(E \Delta T_{t,t+h}(E))}. \end{aligned}$$

Since $(G(t+h, x) - G(t, x))/h$ is L -Lipschitz in x for some $L > 0$, and $\operatorname{div} n_\varphi^E$ is bounded by $4/\rho^2$ in $\{|d_\varphi^E| \leq \rho/2\}$, we deduce

$$\begin{aligned} (20) \quad & \left\| \operatorname{div} n_\varphi^{T_{t,t+h}(E)} + \frac{1}{h}(G(t+h, \cdot) - G(t, \cdot)) \right\|_{L^\infty(\partial T_{t,t+h}(E))} \\ & \leq \left\| \operatorname{div} n_\varphi^E + \frac{1}{h}(G(t+h, \cdot) - G(t, \cdot)) \right\|_{L^\infty(\partial E)} + \frac{4h}{\rho^2} \left(C + \frac{2}{\rho}\right) \\ & \leq \left\| \operatorname{div} n_\varphi^E + \frac{1}{h}(G(t, \cdot) - G(t-h, \cdot)) \right\|_{L^\infty(\partial E)} + h \left(L + \frac{4}{\rho^2}\right) \left(C + \frac{2}{\rho}\right), \end{aligned}$$

provided h is small enough (depending on ρ, L, C). Eventually, it follows that the curvature of $\partial T_{t,t+h}(E)$ (since $d = 2$, the total and mean curvature coincide) has a global estimate $1/\rho + 2C + O(h)$, and one will deduce from Lemma 4 that for h small enough, this new set also satisfies the $\rho'W_\varphi$ -condition, with $\rho' = \rho/(1 + (2C + O(h))\rho) > 0$, provided the assumptions of the lemma are fulfilled.

We now consider E_0, R as in Theorem 1, and let for $h > 0$ and any $n \geq 1$, $E_n^h = T_{(n-1)h, nh}(E_0)$. We also define $E^h(t) = E_{[t/h]}^h$ for $t \geq 0$. A first observation is that if $x \in (E_0)_R$, $x + RW_\varphi \subset E_0$ so that if $r(t)$ solves $\dot{r} = -(1/r + C)$ with $r(0) = R$, for any $\eta > 0$ (small), $x + (r(t) - \eta)W_\varphi \subset E^h(t)$ for h small enough, as long as $r(t) \geq \eta$. The function $r(t)$ solves $r(t) - R - \ln\left(\frac{1+Cr(t)}{1+CR}\right)/C = -Ct$, and given $\delta \in (0, R)$ (which will be precised later on), there exists $T_1(R, C, \delta)$ such that if $t \leq T_1$ and $h > 0$ is small enough,

$$(21) \quad (E_0)_\delta \subset E^h(t) \subset (E_0)^\delta.$$

We let $U = \{|d_\varphi^E| \leq \delta\}$.

Letting $E_1^h = T_{0,h}(E_0)$, we deduce from (20) that if $h < R^2/36$ is small enough,

$$A_1^h := \left\| \operatorname{div} n_\varphi^{E_1^h} + \frac{1}{h}(G(h, \cdot) - G(0, \cdot)) \right\|_{L^\infty(\partial E_1^h)} \leq \frac{1}{R} + C + \frac{1}{4} \left(C + \frac{2}{R}\right) =: M_1.$$

For $n \geq 1$, we then define iteratively the sets $E_{n+1}^h = T_{nh, (n+1)h}(E_n^h)$ and let

$$A_{n+1}^h := \left\| \operatorname{div} n_{\varphi}^{E_{n+1}^h} + \frac{1}{h} (G((n+1)h, \cdot) - G(nh, \cdot)) \right\|_{L^\infty(\partial E_{n+1}^h)}.$$

Let now $R_1 = (2M_1 + C)^{-1}$. As long as $A_n \geq 2M_1$, one can deduce from Lemma 4, using (21) and provided we had chosen $\delta < R_1/2$, that E_{n+1}^h satisfies the $R_1 W_\varphi$ -condition, so that (20) holds (with $E = E_n^h$, $\rho = R_1$) and

$$A_{n+1}^h \leq A_n^h + h \left(L + \frac{4}{R_1^2} \right) \left(C + \frac{2}{R_1} \right).$$

By induction, we deduce that (letting $B = (L + 4/R_1^2)(C + 2/R_1)$) $A_{n+1}^h \leq M_1 + (n+1)hB$ as long as $nh \leq \min\{T_1, M_1/B\} := T > 0$.

We observe that since $\delta < R_1/2$, as long as $nh \leq T$, not only $\partial E_n^h \subset U$, but all the signed distance functions to the boundaries of E_n^h are in $C^{1,1}(U)$. Notice that T and the width δ of the strip U depend only on R, C, L .

Step 1.b.: The case $G_2 = 0$. We now show that we can obtain a similar control in case of a space independent forcing term, which can be the derivative of a continuous function G_2 (a relevant example is a Brownian forcing). In that case, we can consider the algorithm from a different point of view: given the set E , we first consider the set E' with signed distance function $d_{\varphi}^{E'} := d_{\varphi}^E(x) + G(s) - G(t)$, then, we apply to this set E' the algorithm with $G \equiv 0$, that is, we solve (7) for $E = E'$ and $G = 0$:

$$\min_{w \in L^2(B_R)} \int_{B_R} \varphi^\circ(Dw) + \frac{1}{2(s-t)} \int_{B_R} (w(x) - d_{\varphi}^{E'}(x))^2 dx,$$

and then let $E'' = \{w < 0\}$. It is clear that this is equivalent to the original algorithm, so that $E'' = T_{t,s}(E)$.

Assume in addition that E has an inner $r_i W_\varphi$ -condition and a outer $r_o W_\varphi$ condition, for some radii $r_i, r_o > 0$. If $(s-t)$ is small enough, then E' has the inner $r'_i W_\varphi$ -condition and outer $r'_o W_\varphi$ condition with $r'_i = r_i - G(s) + G(t)$ and $r'_o = r_o + G(s) - G(t)$. In particular, $d_{\varphi}^{E'} = d_{\varphi}^E(x) + G(s) - G(t)$ is locally $C^{1,1}$ in the strip $\{-r'_i < d_{\varphi}^{E'} < r'_o\}$ and the surface $\partial E'$ has a curvature which satisfies a.e.

$$(22) \quad -\frac{1}{r'_o} \leq \operatorname{div} n_{\varphi}^{E'} \leq \frac{1}{r'_i}.$$

As before, from (17) we have that, if $h = s - t$ is small enough, then

$$(23) \quad |w(x) - d_{\varphi}^{E'}(x)| \leq \frac{2h}{\min\{r'_i, r'_o\}},$$

showing that the boundary of $T_{t,t+h}(E)$ remains close to the boundary of E' (provided r'_i, r'_o are controlled from below).

From (13) (with $G = 0, E = E'$) and (22), (23), we obtain that if h is small enough,

$$-\frac{1}{r'_o} - \frac{2h}{(r'_o)^2 \min\{r'_i, r'_o\}} \leq \operatorname{div} n_{\varphi}^{T_{t,t+h}(E)} \leq \frac{1}{r'_i} + \frac{2h}{(r'_i)^2 \min\{r'_i, r'_o\}},$$

and in particular we can deduce from Lemma 4 and Remark 4 that $T_h(E)$ satisfies the inner $r''_i W_\varphi$ and outer $r''_o W_\varphi$ -conditions with

$$r''_i \geq r'_i - \frac{ch}{r'_i}, \quad r''_o \geq r'_o - \frac{ch}{r'_o},$$

for some constant $c > 0$.

As in the previous step, we now consider E_0, R as in Theorem 1, we let $E_0^h = E_0$ and define for each $n \geq 0$, $E_{n+1}^h := T_{nh, (n+1)h}(E_n^h)$. Let $r_o^0 = r_i^0 = R$. The previous analysis shows that E_1^h has the inner $r_i^1 W_\varphi$ and the outer $r_o^1 W_\varphi$ -conditions with

$$r_i^1 \geq r_i^0 - G(h) + G(0) - \frac{ch}{R}, \quad r_o^1 \geq r_o^0 + G(h) - G(0) - \frac{ch}{R},$$

provided $|G(h) - G(0)| \leq R/2$ (for some constant $c > 0$). Now, assuming that n is such that

$$r_i^n \geq r_i^0 - G(nh) + G(0) - \frac{cnh}{R}, \quad r_o^n \geq r_o^0 + G(nh) - G(0) - \frac{cnh}{R},$$

we deduce that

$$r_i^{n+1} \geq r_i^0 - G((n+1)h) + G(0) - \frac{c(n+1)h}{R}, \quad r_o^{n+1} \geq r_o^0 + G((n+1)h) - G(0) - \frac{c(n+1)h}{R},$$

as long as $|G((n+1)h) - G(0)| + c(n+1)h/R \leq R/2$. Define T such that $\max_{0 \leq t \leq T} |G(t) - G(0)| + ct/R \leq R/4$, and let $U = \{|d_\varphi^{E_0}| < R/4\}$: then, on one hand, $\partial E_n^h \subset U$ for all $n \geq 0$ with $nh \leq T$, on the other hand, E_n^h satisfies the $(R/2)W_\varphi$ -condition, so that $d_\varphi^{E_n^h} \in C^{1,1}(U)$. Again, U and T depend only on G and R .

Step 3: Conclusion. For $t \in [0, T]$ and h small, we let $E_h(t) = E_{[t/h]}^h$, $d_h(t, x) = d_\varphi^{E_h(t)}(x)$, and we now send $h \rightarrow 0$. Since $d_h - G$ is uniformly Lipschitz in $[0, T] \times U$ (in time, in fact, we have $|d_h(t, x) - G(t, x) - d_h(s, x) + G(s, x)| \leq c|t - s|$ if $|t - s| \geq h$, for some constant c), up to a subsequence (h_k) it converges uniformly to some d with $d - G \in \text{Lip}([0, T] \times U)$, moreover, at each $t > 0$, $E_{h_k}(t)$ converges (Hausdorff) to a set $E(t)$ with $d(t, x) = d_\varphi^{E(t)}(x)$. Let us establish (18).

For $n \leq T/h - 1$ and $x \in \partial E_{n+1}^h$, by definition of the scheme we have

$$-d_\varphi^{E_n^h}(x) - h \operatorname{div} n_\varphi^{E_{n+1}^h}(x) - G(t+h, x) + G(t, x) = 0.$$

As $(G(t+h, \cdot) - G(t, \cdot))/h$ is L -Lipschitz in U , there holds

$$\left| (G(t+h, x) - G(t, x)) - (G(t+h, \Pi_{\partial E_{n+1}^h}(x)) - G(t, \Pi_{\partial E_{n+1}^h}(x))) \right| \leq Ch |d_\varphi^{E_{n+1}^h}(x)|$$

where C depends only on L and φ , where we set

$$\Pi_{\partial E_{n+1}^h}(x) = x - d_\varphi^{E_{n+1}^h}(x) n_\varphi^{E_{n+1}^h}(x).$$

Choose now $x \in U$ such that $d_\varphi^{E_{n+1}^h}(x) \geq 0$. In this case, it follows that

$$d_\varphi^{E_n^h}(x) - d_\varphi^{E_n^h}(\Pi_{\partial E_{n+1}^h}(x)) \leq \varphi(x - \Pi_{\partial E_{n+1}^h}(x)) = d_\varphi^{E_{n+1}^h}(x).$$

Hence,

$$\begin{aligned} & d_\varphi^{E_{n+1}^h}(x) - d_\varphi^{E_n^h}(x) - h \operatorname{div} n_\varphi^{E_{n+1}^h}(x) \\ & \geq -d_\varphi^{E_n^h}(\Pi_{\partial E_{n+1}^h}(x)) - h \operatorname{div} n_\varphi^{E_{n+1}^h}(\Pi_{\partial E_{n+1}^h}(x)) + O\left(h |d_\varphi^{E_{n+1}^h}(x)|\right) \\ & = G(t+h, \Pi_{\partial E_{n+1}^h}(x)) - G(t, \Pi_{\partial E_{n+1}^h}(x)) + O\left(h |d_\varphi^{E_{n+1}^h}(x)|\right) \\ & = G(t+h, x) - G(t, x) + O\left(h |d_\varphi^{E_{n+1}^h}(x)|\right). \end{aligned}$$

Dividing by h and letting $h \rightarrow 0^+$, we then get

$$\frac{\partial(d_\varphi^E - G)}{\partial t}(t, x) - \operatorname{div} \nabla \varphi^\circ(\nabla d_\varphi^E)(t, x) \geq O(|d_\varphi^E(t, x)|) \quad (t, x) \in U \times [0, T] \cap \{d_\varphi^E(t, x) > 0\},$$

which implies

$$\frac{\partial(d_\varphi^E - G)}{\partial t}(t, x) - \operatorname{div} \nabla \varphi^\circ(\nabla d_\varphi^E)(t, x) \geq O(|d_\varphi^E(t, x)|) \quad (t, x) \in U \times [0, T].$$

By taking $x \in U$ such that $d_\varphi^{E_h}(x) \leq 0$, reasoning as above we get

$$\frac{\partial d_\varphi^E}{\partial t}(t, x) - \operatorname{div} \nabla \varphi^\circ(\nabla d_\varphi^E)(t, x) - g(t, x) \leq O(|d_\varphi^E(t, x)|) \quad (t, x) \in U \times [0, T],$$

thus obtaining (18). Q.E.D.

Remark 5. When $\varphi(x) = |x|$ and $G_2 = 0$, the an existence and uniqueness result for φ -regular flows has been proved in [18] in any dimension.

4.2. Uniqueness of φ -regular flows.

We now show uniqueness of the regular evolutions given by Theorem 1.

Theorem 2. *Given an initial set E_0 , the flow of Theorem 1 is unique. More precisely, if two flows E, E' are given, starting from initial sets $E_0 \subseteq E'_0$, then $E(t) \subseteq E'(t)$ for all $t \in [0, \min\{T, T'\}]$ (where T, T' are respectively the time of existence of regular flows starting from E_0 and E'_0).*

The thesis essentially follows from the results in [15]. Indeed, in [15] it is proved a comparison result for strict C^2 sub- and superflows, based again on a consistency result for the scheme defined in Section 3. A strict C^2 subflow is defined a in Theorem 1, except that $d_\varphi^E(t, x)$ is required to be in $C^0([0, T]; C^2(U))$, and (18) is replaced with (for $0 \leq t < s \leq T, x \in U$)

$$(24) \quad d_\varphi^E(s, x) - d_\varphi^E(t, x) - \int_s^t \operatorname{div} \nabla \varphi^\circ(\nabla d_\varphi^E)(\tau, x) d\tau - G(s, x) + G(t, x) \leq -\delta(s - t)$$

for some $\delta > 0$. A superflow will satisfy the reverse inequality, with $-\delta(s - t)$ replaced with $\delta(s - t)$. For technical reasons (in order to make sure, in fact, that the duration time of these flows is independent on δ), we will ask that these flows are defined, in fact, in a tubular neighborhood W of $\bigcup_{0 \leq t \leq T} \partial E(t)$, not necessarily of the form $[0, T] \times U$.

The thesis then follows from the consistency result in [15, Thm. 3.3], once we show the following approximation result.

Lemma 5. *Let $E(t)$ be an evolution as in Theorem 1, starting from a compact set E_0 satisfying the RW_φ -conditions for some $R > 0$. Then, for any $\varepsilon > 0$, there exists $T' > 0$ (depending only on R), a set E'_0 and a strict C^2 subflow $E'(t)$ starting from E'_0 such that for all $t \in [0, T']$, $E(t) \subset \subset E'(t) \subset \{d_\varphi^E \leq \varepsilon\}$.*

Proof. We sketch the proof and refer to [3] for more details.

The idea is to let first $d^\alpha = d_\varphi^E - \alpha t - \alpha/(4\lambda)$, for some small $\alpha > 0$, with $\alpha(T + 1/(4\lambda)) < \varepsilon$. One can then deduce from (18) that, for all $s > t$,

$$\begin{aligned} d^\alpha(s, x) - d^\alpha(t, x) - \int_s^t \operatorname{div} \nabla \varphi^\circ(\nabla d^\alpha)(\tau, x) d\tau - G(s, x) + G(t, x) \\ \leq (s - t)(\lambda \max_{t \leq \tau \leq s} |d_\varphi^E(x, \tau)| - \alpha) + \leq \lambda(s - t) \left(\max_{t \leq \tau \leq s} |d^\alpha(x, \tau)| + \alpha(s - \frac{3}{4}\lambda^{-1}) \right). \end{aligned}$$

Let $T' := \min\{T, 1/(2\lambda)\}$, and let $\beta = \alpha/(8\lambda)$: then if we let $W = \{(t, x) : 0 \leq t \leq T, |d^\alpha(t, x)| < \beta\}$, we deduce that for $(t, x), (s, x) \in W$,

$$d^\alpha(s, x) - d^\alpha(t, x) - \int_s^t \operatorname{div} \nabla \varphi^\circ(\nabla d^\alpha)(\tau, x) d\tau - G(s, x) + G(t, x) \leq -\beta(t - s).$$

Hence $\{d^\alpha \leq 0\}$ is almost a C^2 subflow, except for the fact that it is not C^2 . However, this is not really an issue, as we will now check. Consider indeed a spatial mollifier

$$(25) \quad \eta_r(x) = \frac{1}{r^d} \eta\left(\frac{x}{r}\right)$$

where as usual $\eta \in C_c^\infty(B(0, 1); \mathbb{R}_+)$, $\int_{\mathbb{R}^d} \eta(x) dx = 1$. Let $d_r^\alpha = \eta_r * d^\alpha$ (for r small). Observing, as before, that $G(s, x) - G(t, x)$ is $(s - t)L$ -Lipschitz in x , one has $|\eta_r * (G(s, \cdot) - G(t, \cdot))(x) - (G(s, x) - G(t, x))| \leq (s - t)Lr$. Hence, the level set 0 of d_r^α will be a strict C^2 subflow, for r small enough, if we can check that the difference

$$(26) \quad \eta_r * (\operatorname{div} \nabla \varphi^\circ(\nabla d^\alpha)(\tau, \cdot))(x) - \operatorname{div} \nabla \varphi^\circ(\eta_r * \nabla d^\alpha)(\tau, x)$$

can be made arbitrarily small for r small enough and any $(\tau, x) \in W$ (possibly reducing slightly the width of W). Now, for $(\tau, x) \in W$,

$$\eta_r * (\operatorname{div} \nabla \varphi^\circ(\nabla d^\alpha)(\tau, \cdot))(x) = \int_{B(0, r)} \eta_r(z) D^2 \varphi^\circ(\nabla d^\alpha(\tau, x - z)) : D^2 d^\alpha(\tau, x - z) dz$$

while

$$\operatorname{div} \nabla \varphi^\circ(\eta_r * \nabla d^\alpha)(\tau, x) = \int_{B(0, r)} \eta_r(z) D^2 \varphi^\circ((\eta_r * \nabla d^\alpha(\tau, \cdot))(x)) : D^2 d^\alpha(\tau, x - z) dz.$$

The difference in (26) is therefore

$$\int_{B(0, r)} \eta_r(z) (D^2 \varphi^\circ(\nabla d^\alpha(\tau, x - z)) - D^2 \varphi^\circ((\eta_r * \nabla d^\alpha(\tau, \cdot))(x))) : D^2 d^\alpha(\tau, x - z) dz.$$

Now, since $D^2 \varphi^\circ$ is at least continuous (uniformly in $\{\varphi^\circ(\xi) \geq 1/2\}$), $\varphi^\circ(\nabla d^\alpha) = 1$ a.e. in W , while $D^2 d^\alpha$ is globally bounded (and ∇d^α uniformly Lipschitz), this difference can be made arbitrarily small as $r \rightarrow 0$, and we actually deduce that, in such a case, $E'(t) = \{d_r^\alpha \leq 0\}$ is a strict C^2 -superflow starting from $E'_0 = \{d_\varphi^E \leq \beta\}$, which satisfies the thesis of the Lemma. Q.E.D.

Remark 6. The uniqueness result holds in any dimension $d \geq 2$, with exactly the same proof. It is also not necessary to assume that G_1 or G_2 vanishes.

§5. General anisotropies

An important feature of Theorem 1 is that the existence time, as well as the neighborhood where d_φ^E is $C^{1,1}$, are both independent on the anisotropy, and only depend on the radius R for which E_0 satisfies the RW_φ -condition. This allows us to extend the existence result to general anisotropies, by the approximation argument given in Lemma 1.

Theorem 3. *Assume $G_1 = 0$ or $G_2 = 0$, and let (φ, φ°) be an arbitrary anisotropy. Let $E_0 \subset \mathbb{R}^2$ an initial set with compact boundary, satisfying the RW_φ -condition for some $R > 0$. Then, there exist $T > 0$, and a φ -regular flow $E(t)$ defined on $[0, T]$ and starting from E_0 .*

More precisely, there exist $R' > 0$ and a neighborhood U of $\bigcup_{0 \leq t \leq T} \partial E(t)$ in \mathbb{R}^2 such that the sets $E(t)$ satisfy the $R'W_\varphi$ -condition for all $t \in [0, T]$, the φ -signed distance function $d_\varphi^E(t, x)$ from $\partial E(t)$ belongs to $C^0([0, T]; \operatorname{Lip}(U))$, $(d_\varphi^E - G) \in \operatorname{Lip}([0, T] \times U)$ and

$$(27) \quad \left| \frac{\partial(d_\varphi^E - G)}{\partial t}(t, x) - \operatorname{div} z(t, x) \right| \leq \lambda |d_\varphi^E(t, x)|$$

for a.e. $(t, x) \in [0, T] \times U$, where λ is a positive constant and $z \in L^\infty([0, T] \times U; \mathbb{R}^2)$ is such that $z \in \partial \varphi^\circ(\nabla d_\varphi^E)$ a.e. in $[0, T] \times U$. The time T , the radius R' , and the constant λ , only depend on R and G .

Remark 7. Comparison and uniqueness for such flows has been shown in [11, 7, 14], although the most general result in these references only covers the case of a time-dependent, Lipschitz continuous forcing term $G(t) = G(0) + \int_0^t c(s) ds$, with $c \in L^\infty(0, +\infty)$.

Proof. Let $\varepsilon > 0$ and consider smooth and elliptic anisotropies $(\varphi_\varepsilon, \varphi_\varepsilon^\circ)$, with $\varphi_\varepsilon \geq \varphi$, converging to (φ, φ°) locally uniformly as $\varepsilon \rightarrow 0$. By the approximation result in Lemma 1, we can find a sequence of sets E_ε which satisfy the RW_{φ_ε} -condition, and such that $\partial E_\varepsilon \rightarrow \partial E$ in the Hausdorff sense. For each ε we consider the evolution $E^\varepsilon(t)$ given by Theorem 1, with $0 \leq t \leq T^\varepsilon$. Since the times T^ε and the width of the neighborhoods U^ε depend only on R and G , up to extracting a subsequence we can assume that $\lim_\varepsilon T^\varepsilon = T$ for some $T > 0$, and there exists a neighborhood U of ∂E_0 such that $\mathbb{R}^d \setminus U^\varepsilon$ converges to $\mathbb{R}^d \setminus U$ in the Hausdorff sense, as $\varepsilon \rightarrow 0$. Possibly reducing T and the width of U we can then assume that $T^\varepsilon = T$ and $U^\varepsilon = U$ for all $\varepsilon > 0$.

Letting $W := [0, T] \times U$, and $z_\varepsilon(t, x) = \nabla \varphi_\varepsilon^\circ(\nabla d_{\varphi_\varepsilon}^{E_\varepsilon}(t, x))$, from (18) we get

$$(28) \quad \left| \frac{\partial(d_{\varphi_\varepsilon}^{E_\varepsilon} - G)}{\partial t}(t, x) - \operatorname{div} z_\varepsilon(t, x) \right| \leq \lambda |d_{\varphi_\varepsilon}^{E_\varepsilon}(t, x)|$$

for a.e. $(t, x) \in W$, where the constant λ depends only on R and G .

As $d_{\varphi_\varepsilon}^{E_\varepsilon} - G$ are uniformly Lipschitz in (t, x) , up to a subsequence we can assume that the functions $d_{\varphi_\varepsilon}^{E_\varepsilon}$ converge uniformly in any compact subset of W to a function d_φ^E , such that for all $t \in [0, T]$ $d_\varphi^E(t, \cdot)$ is the signed φ -distance function to the boundary of $E(t) := \{x : d_\varphi^E(t, x) \leq 0\}$. Moreover, $E(0) = E_0$, $E(t)$ is the Hausdorff limit of $E_\varepsilon(t)$ for each $t \in [0, T]$, and satisfies the $R'W_\varphi$ -condition, with $R' = \lim_\varepsilon R'_\varepsilon$.

Up to a subsequence we can also assume that there exists $z \in L^\infty(W)$ with $z_\varepsilon(t, x) \xrightarrow{*} z(t, x)$, $\operatorname{div} z_\varepsilon \xrightarrow{*} \operatorname{div} z$ and $\partial_t(d_{\varphi_\varepsilon}^{E_\varepsilon} - G) \xrightarrow{*} \partial_t(d_\varphi^E - G)$ in $L^\infty(W)$, so that (27) holds a.e. in W .

It remains to check that $z(t, x) \in \partial \varphi^\circ(\nabla d_\varphi^E(t, x))$. Since by construction $z(t, x) \in W_\varphi$ for a.e. $(t, x) \in W$, it is enough to show that

$$(29) \quad z \cdot \nabla d_\varphi^E = \varphi^\circ(\nabla d_\varphi^E) = 1$$

a.e. in W . Recalling that $z_\varepsilon \cdot \nabla d_{\varphi_\varepsilon}^{E_\varepsilon} = \varphi_\varepsilon^\circ(\nabla d_{\varphi_\varepsilon}^{E_\varepsilon}) = 1$ a.e. in W and letting $\psi \in C_c^\infty(W)$, we have

$$\int_W \psi \, dxdt = \int_W \psi (z_\varepsilon \cdot \nabla d_{\varphi_\varepsilon}^{E_\varepsilon}) \, dxdt = - \int_W d_{\varphi_\varepsilon}^{E_\varepsilon} (z_\varepsilon \cdot \nabla \psi + \psi \operatorname{div} z_\varepsilon) \, dxdt.$$

Passing to the limit in the right-hand side we then get

$$\int_W \psi \, dxdt = - \int_W d_\varphi^E (z \cdot \nabla \psi + \psi \operatorname{div} z) \, dxdt = \int_W \psi (z \cdot \nabla d_\varphi^E) \, dxdt$$

which gives (29). Q.E.D.

References

- [1] F. J. Almgren and J. E. Taylor. Flat flow is motion by crystalline curvature for curves with crystalline energies. *J. Diff. Geom.*, 42:1–22, 1995.
- [2] F. J. Almgren, J. E. Taylor and L.-H. Wang. Curvature-driven flows: a variational approach *SIAM J. Control Optim.*, 31(2):387–438, 1993.
- [3] L. Almeida, A. Chambolle and M. Novaga. Mean curvature flow with obstacles. *Annales IHP - Analyse Nonlineaire*, to appear.
- [4] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford Mathematical Monographs, 2000.
- [5] S. Angenent. Parabolic equations for curves on surfaces I. Curves with p-integrable curvature. *Ann. of Math. (2)*, 132(3):451–483, 1990.
- [6] G. Bellettini, V. Caselles, A. Chambolle and M. Novaga. Crystalline mean curvature flow of convex sets. *Arch. Rat. Mech. Anal.*, 179(1):109–152, 2006.

- [7] G. Bellettini, V. Caselles, A. Chambolle and M. Novaga. The volume preserving crystalline mean curvature flow of convex sets in \mathbb{R}^N . *Journal de mathématiques pures et appliquées*, 92(5):499–527, 2009.
- [8] G. Bellettini, R. Gogione and M. Novaga, Approximation to driven motion by crystalline curvature in two dimensions, *Adv. Math. Sci. and Appl.*, 10:467–493, 2000.
- [9] G. Bellettini, M. Novaga and M. Paolini. On a crystalline variational problem I. First variation and global L^∞ regularity. *Arch. Ration. Mech. Anal.* 157(3), 165–191, 2001.
- [10] G. Bellettini, M. Novaga, and M. Paolini. On a crystalline variational problem. II. BV regularity and structure of minimizers on facets. *Arch. Ration. Mech. Anal.* 157(3), 193–217, 2001.
- [11] G. Bellettini and M. Novaga. Approximation and comparison for non-smooth anisotropic motion by mean curvature in \mathbb{R}^N . *Math. Mod. Meth. Appl. Sc.*, **10** (2000), 1–10.
- [12] J.W. Cahn and D.W. Hoffman. A vector thermodynamics for anisotropic interfaces. 1. Fundamentals and applications to plane surface junctions. *Surface Sci.*, 31:368–388, 1972.
- [13] A. Chambolle. An algorithm for mean curvature motion. *Interfaces Free Bound.*, 6:195–218, 2004.
- [14] A. Chambolle and M. Novaga. Convergence of an algorithm for the anisotropic and crystalline mean curvature flow. *SIAM J. Math. Anal.*, 37(6):1978–1987 (electronic), 2006.
- [15] A. Chambolle and M. Novaga. Implicit time discretization of the mean curvature flow with a discontinuous forcing term. *Interfaces Free Bound.*, 10:283–300, 2008.
- [16] M. G. Crandall, H. Ishii and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Am. Math. Soc.*, 27:1–67, 1992.
- [17] Y. G. Chen, Y. Giga and S. Goto. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. *J. Differential Geom.*, 33(3):749–786, 1991.
- [18] N. Dirr, S. Luckhaus, M. Novaga. A stochastic selection principle in case of fattening for curvature flow. *Calc. Var. PDE*, 13(4):405–425, 2001.
- [19] L. C. Evans Convergence of an algorithm for mean curvature motion. *Indiana Univ. Math. J.*, 42:533–557, 1993.
- [20] L. C. Evans and R. F. Gariepy. *Measure Theory and Fine Properties of Functions*. Studies in Advanced Math., CRC Press, Ann Harbor, 1992.
- [21] L. C. Evans and J. Spruck. Motion of level sets by mean curvature. II. *Trans. Amer. Math. Soc.*, 330:321–332, 1992.
- [22] M. Gage and R. Hamilton. The heat equation shrinking convex plane curves. *J. Differential Geom.*, 23:69–95, 1986.
- [23] M.-H. Giga, Y. Giga and P. Rybka, A comparison principle for singular diffusion equation with spatially inhomogeneous driving force for graphs. (preprint # 981, Hokkaido University, 2011).
- [24] Y. Giga, P. Górka and P. Rybka, Evolution of regular bent rectangles by the driven crystalline curvature flow in the plane with a non-uniform forcing term. (preprint # 993, Hokkaido University, 2011).
- [25] Y. Giga and M. E. Gurtin. A comparison theorem for crystalline evolutions in the plane. *Quarterly of Applied Mathematics*, LIV:727–737, 1996.
- [26] P.-L. Lions and P. E. Souganidis. Fully nonlinear stochastic partial differential equations. *C.R. Acad. Sci. - Series I - Mathematics* 326(9):1085–1092, 1998.
- [27] S. Luckhaus and Sturzenecker, Implicit time discretization for the mean curvature flow equation. *Calc. Var.* **3** (1995), 253–271.
- [28] J. E. Taylor. Crystalline variational problems. *Bull. Amer. Math. Soc. (N.S.)*, 84:568–588, 1978.
- [29] J. E. Taylor, Motion of curves by crystalline curvature, including triple junctions and boundary points, *Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990)*, 417–438, Proc. Sympos. Pure Math., 54, Part 1, Amer. Math. Soc., Providence, 1993

CMAP, Ecole Polytechnique, CNRS, 91128 Palaiseau, France,
antonin.chambolle@cmap.polytechnique.fr

Dipartimento di Matematica, Università di Pisa, Largo Pontecorvo 5, 56127 Pisa, Italy,
novaga@dm.unipi.it